



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Health conditions in La Guaira remain good. No unusual mortality is observed among rats. One fatal case of plague occurred at Caracas July 2.

ZANZIBAR.

Report from Zanzibar—Plague present.

The following is received from the Department of State under date of July 26:

Cablegram from Zanzibar reports "One case plague has appeared."

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

AFRICA—*Lourenço Marquez*.—Month of May, 1909. Estimated population, 10,000. Total number of deaths, 50, corresponding to an annual death rate of 60 per 1,000 of population, including 12 from tuberculosis.

ALGERIA—*Bona*.—Month of June, 1909. Estimated population, 42,934. Total number of deaths, 94, corresponding to an annual death rate of 26.28 per 1,000 of population, including smallpox 7, enteric fever 2, typhus fever 10, scarlet fever 1, and 15 from tuberculosis.

ARGENTINA—*Buenos Aires*.—Month of April, 1909. Estimated population, 1,201,722. Total number of deaths, 1,360, corresponding to an annual death rate of 13.2 per 1,000 of population, including enteric fever 37, scarlet fever 3, whooping cough 1, smallpox 6, measles 2, diphtheria 6, leprosy 1, and 157 from tuberculosis.

BRAZIL—*Bahia*.—Month of March, 1909. Estimated population, 286,000. Total number of deaths, 465, corresponding to an annual death rate of 19.15 per 1,000 of population, including yellow fever 24, plague 1, smallpox 8, beriberi 4, diphtheria 1, enteric fever 2, and 55 from pulmonary tuberculosis.

São Paulo, Santos, and Campinas.—Two weeks ended June 13, 1909. Estimated population, 300,000. Total number of deaths, 314, corresponding to an annual death rate of 27.04 per 1,000 of population, including leprosy 1, enteric fever 2, smallpox 3, scarlet fever 1, diphtheria 1, and 18 from tuberculosis.

CHILE—*Punta Arenas*.—Month of May, 1909. Estimated population, 12,000. Total number of deaths, 23, corresponding to an annual death rate of 21.8 per 1,000 of population, including 5 from tuberculosis.

FRANCE—*Cherbourg*.—Month of June, 1909. Estimated population, 43,948. Total number of deaths, 84, corresponding to an annual death rate of 22.8 per 1,000 of population, including enteric fever 3, and 19 from tuberculosis.